

Referred Publications

1. Lawson, S. J., Galbally, I. E., Powell, J. C., Keywood, M. D., Molloy, S. B., Cheng M., and Selleck, P. W. (2011). The effect of proximity to major roads on indoor air quality in typical Australian dwellings. *Atmospheric Environment*, in press.
2. Galbally, I. E., Keywood, M. D., Powell, J. C., Lawson, S. J., Cheng M., Dunne, R., Gillett, R. W., Molloy, S. B., Selleck, P. W., Ward, J., and Reisen, F. (2011). An overview of the CSIRO 2008-2009 indoor air quality study. *Clean Air and Environmental Quality*, in press.
3. Radhi, M., Box, M. A., Box, G. P., Mitchell, R. M., Cohen, D. D., Stelcer, E., and Keywood, M. D. (2010a). Optical, physical and chemical characteristics of Australian continental aerosols: results from a field experiment. *Atmospheric Chemistry and Physics*, 10(13), 5925-5942.
4. Gabric, A. J., Cropp, R. A., McTainsh, G. H., Johnston, B. M., Butler, H., Tilbrook, B., and Keywood, M. (2010). Australian dust storms in 2002-2003 and their impact on Southern Ocean biogeochemistry. *Global Biogeochemical Cycles*, 24.
5. Radhi, M., Box, M. A., Box, G. P., Mitchell, R. M., Cohen, D. D., Stelcer, E., and Keywood, M. D. (2010b). Size-resolved mass and chemical properties of dust aerosols from Australia's Lake Eyre Basin. *Atmospheric Environment*, 44(29), 3519-3528.
6. Rotstayn, L. D., Keywood, M. D., Forgan, B. W., Gabric, A. J., Galbally, I. E., Gras, J. L., et al. (2009). Possible impacts of anthropogenic and natural aerosols on Australian climate: a review. *International Journal of Climatology*, 29(4), 461-479.
7. Iinuma, Y., Boge, O., Keywood, M., Gnauk, T., & Herrmann, H. (2009). Diaterpeneic Acid Acetate and Diaterpenylic Acid Acetate: Atmospheric Tracers for Secondary Organic Aerosol Formation from 1,8-Cineole Oxidation. *Environmental Science & Technology*, 43(2), 280-285.
8. O'Toole, J., Keywood, M., Sinclair, M., & Leder, K. (2009). Risk in the mist? Deriving data to quantify microbial health risks associated with aerosol generation by water-efficient devices during typical domestic water-using activities. *Water Science and Technology*, 60(11), 2913-2920.
9. Cainey, J. M., Keywood, M., Bigg, E. K., Grose, M. R., Gillett, R. W., & Meyer, M. (2007). Flux chamber study of particle formation from *Durvillaea potatorum*. *Environmental Chemistry*, 4(3), 151-154.
10. Cainey, J. M., Keywood, M., Grose, M. R., Krummel, P., Galbally, I. E., Johnston, P., et al. (2007). Precursors to Particles (P2P) at Cape Grim 2006: campaign overview. *Environmental Chemistry*, 4(3), 143-150.
11. Keywood, M. D. (2007). Aerosol composition at Cape Grim : an evaluation of PM10 sampling program and baseline event switches. *Baseline Atmospheric Program Australia 2005-2006. 2005-2006 ed. J. M. Cainey, N. Derek, and P. B. Krummel (editors). Melbourne: Australian Bureau of Meteorology and CSIRO Marine and Atmospheric Research. p. 31-36.*
12. Lee, A., Goldstein, A. H., Keywood, M. D., Gao, S., Varutbangkul, V., Bahreini, R., et al. (2006). Gas-phase products and secondary aerosol yields from the ozonolysis of ten different terpenes. *Journal of Geophysical Research-Atmospheres*, 111(D7).
13. Luhar, A. K., Galbally, I. E., & Keywood, M. (2006). Modelling PM10 concentrations and carrying capacity associated with woodheater emissions in Launceston, Tasmania. *Atmospheric Environment*, 40(29), 5543-5557.
14. Varutbangkul, V., Brechtel, F. J., Bahreini, R., Ng, N. L., Keywood, M. D., Kroll, J. H., et al. (2006). Hygroscopicity of secondary organic aerosols formed by oxidation of cycloalkenes, monoterpenes, sesquiterpenes, and related compounds. *Atmospheric Chemistry and Physics*, 6, 2367-2388.
15. Bahreini, R., Keywood, M. D., Ng, N. L., Varutbangkul, V., Gao, S., Flagan, R. C., et al. (2005). Measurements of secondary organic aerosol from oxidation of cycloalkenes, terpenes, and m-xylene using an Aerodyne aerosol mass spectrometer. *Environmental Science & Technology*, 39(15), 5674-5688.
16. Ng, N. L., Kroll, J. H., Keywood, M. D., Bahreini, R., Varutbangkul, V., Flagan, R. C., et al. (2006). Contribution of first- versus second-generation products to secondary organic aerosols formed in the oxidation of biogenic hydrocarbons. *Environmental Science & Technology*, 40(7), 2283-2297.

17. Keywood, M. D., Kroll, J. H., Varutbangkul, V., Bahreini, R., Flagan, R. C., & Seinfeld, J. H. (2004). Secondary organic aerosol formation from cyclohexene ozonolysis: Effect of OH scavenger and the role of radical chemistry. *Environmental Science & Technology*, 38(12), 3343-3350.
18. Keywood, M. D., Varutbangkul, V., Bahreini, R., Flagan, R. C., & Seinfeld, J. H. (2004). Secondary organic aerosol formation from the ozonolysis of cycloalkenes and related compounds. *Environmental Science & Technology*, 38(15), 4157-4164.
19. Schauer, J. J., Mader, B. T., Deminter, J. T., Heidemann, G., Bae, M. S., Seinfeld, J. H., et al. (2003). ACE-Asia intercomparison of a thermal-optical method for the determination of particle-phase organic and elemental carbon. *Environmental Science & Technology*, 37(5), 993-1001.
20. Gao, S., Keywood, M., Ng, N. L., Surratt, J., Varutbangkul, V., Bahreini, R., et al. (2004). Low-molecular-weight and oligomeric components in secondary organic aerosol from the ozonolysis of cycloalkenes and alpha-pinene. *Journal of Physical Chemistry A*, 108(46), 10147-10164.
21. Gao, S., Ng, N. L., Keywood, M., Varutbangkul, V., Bahreini, R., Nenes, A., et al. (2004). Particle phase acidity and oligomer formation in secondary organic aerosol. *Environmental Science & Technology*, 38(24), 6582-6589.
22. Keywood, M. D. (2003). Aerosol chemistry at Cape Grim. *Baseline Atmospheric Program Australia. 1999-2000 ed. N. W. Tindale, N. Derek, and P. J. Fraser (editors). Melbourne: Bureau of Meteorology and CSIRO Atmospheric Research. p. 31-41.*
23. Keywood, M. D., Ayers, G. P., Gras, J. L., Boers, R., & Leong, C. P. (2003). Haze in the Klang Valley of Malaysia. *Atmospheric Chemistry and Physics*, 3, 591-605.
24. Gras, J. L., Keywood, M. D., & Ayers, G. P. (2001). Factors controlling winter-time aerosol light scattering in Launceston, Tasmania. *Atmospheric Environment*, 35(10), 1881-1889.
25. O'Leary, R., Ayers, G. P., Bridgman, H. A., and Keywood, M. D (2002) Indoor/outdoor air pollution relationships at a household level in SE Australia: a pilot study. *Clean Air*, 36 (2), 25-31
26. Keywood, M. D., Ayers, G. P. and Gras, J. L. (2000) Testing a low-cost aerosol sampler, *Clean Air*, 43, 38-42.
27. Keywood, M. D., Ayers, G. P., Gras, J. L., Gillett, R. W., & Cohen, D. (1999). An evaluation of PM10 and PM2.5 size selective inlet performance using ambient aerosol. *Aerosol Science and Technology*, 30(4), 401-407.
28. Keywood, M. D., Ayers, G. P., Gras, J. L., Gillett, R. W., & Cohen, D. D. (1999). Relationships between size segregated mass concentration data and ultrafine particle number concentrations in urban areas. *Atmospheric Environment*, 33(18), 2907-2913.
29. Keywood, M. D., Ayers, G. P., Gras, J. L., Gillett, R. W., & Cohen, D. D. (2000). Size distribution and sources of aerosol in Launceston, Australia, during winter 1997. *Journal of the Air & Waste Management Association*, 50(3), 418-427.
30. Ayers, G. P., Keywood, M. D., & Gras, J. L. (1999). TEOM vs. manual gravimetric methods for determination of PM2.5 aerosol mass concentrations. *Atmospheric Environment*, 33(22), 3717-3721.
31. Ayers, G. P., Keywood, M. D., Gillett, R., Manins, P. C., Malfroy, H., & Bardsley, T. (1998). Validation of passive diffusion samplers for SO2 and NO2. *Atmospheric Environment*, 32(20), 3587-3592.
32. Ayers, G. P., Selleck, P. W., Gillett, R. W., & Keywood, M. D. (1998). Determination of nicotine in water by gradient ion chromatography. *Journal of Chromatography A*, 824(2), 241-245.
33. Keywood, M. D., Ayers, G. P., Gras, J. L. and Cohen, D. D. (1998) The Use of a Micro-orifice Uniform Deposit Impactor (MOUDI) to investigate relationships between PM10, PM2.5, PM1 and Ultrafine Particles in Urban Australia. *Journal of Aerosol Science*, 29 S1, 99-100.
34. Keywood, M. D., Beer, T., Ayers, G. P, Gillett, R. W. and Kreibich, H. (1998) The use of passive gas samplers to monitor personal exposure. *Clean Air*, 32, 32-36. Ayers G.P., Selleck P.W., Gillett R.W. and Keywood M.D., (1998) Determination of nicotine in water by gradient ion chromatography, *Journal of Chromatography A*, 824, 241-245.

35. Keywood, M. D., Fifield, L. K., Chivas, A. R., & Cresswell, R. G. (1998). Fallout of chlorine 36 to the Earth's surface in the southern hemisphere. *Journal of Geophysical Research-Atmospheres*, 103(D7), 8281-8286.
36. Keywood, M. D., Chivas, A. R., Fifield, L. K., Cresswell, R. G., & Ayers, G. P. (1997). The accession of chloride to the western half of the Australian continent. *Australian Journal of Soil Research*, 35(5), 1177-1189.
37. Simmons, S. F., Keywood, M., Scott, B. J., & Keam, R. F. (1993). IRREVERSIBLE CHANGE OF THE ROTOMAHANA-WAIMANGU HYDROTHERMAL SYSTEM (NEW-ZEALAND) AS A CONSEQUENCE OF A VOLCANIC-ERUPTION. *Geology*, 21(7), 643-646.
38. Keywood, M. D., Rodgers, K. A., Nicholson, K. A (1992) Catalogue of Active Thermal Features in the Waimangu Valley, 1990. *New Zealand Volcanological Record* 19, 29-51.